
Directions:

- 1. Section: Math251 007
- 2. Write your name with one character in each box below.
- 3. Show all work. No credit for answers without work.
- 4. This assessment is closed book and closed notes. You may not use electronic devices, including calculators, laptops, and cell phones.

Academic Integrity Statement: I will complete this work on my own without assistance, knowing or otherwise, from anyone or anything other than the instructor. I will not use any electronic equipment or notes (except as permitted by an existing official, WVU-authorized accommodation).

S	lignatı	ire:				

- 1. Evaluate the following.
 - (a) [4 points] $\iint_R \sin(y^2) dA$ where R is the triangular region bounded by y = x, y = 2x, and y = 1.




$$\iint_{R} \sin(y^{2}) dA = \iint_{D} \iint_{\frac{1}{2}}^{\frac{1}{2}} \sin(y^{2}) dx dy = \iint_{D} \left[\sin(y^{2}) \iint_{\frac{1}{2}} 1 dx \right] dy$$

$$= \iint_{D} \sin(y^{2}) \left[x \right]_{x=\frac{1}{2}}^{x=y} dy = \iint_{D} \left[\sin(y^{2}) \right] \left[y - \frac{1}{2} \right] dy = \frac{1}{2} \iint_{D} y \sin(y^{2}) dy \qquad u = y^{2} du = 2y dy$$

$$= \frac{1}{4} \iint_{D} \sin(y^{2}) \left[2y dy \right] = \frac{1}{4} \iint_{D^{2}}^{2} \sin(u) du = \frac{1}{4} \left(-\cos(u) \right]_{u=0}^{1} = \frac{1}{4} \left(-\cos(u) + \cos(u) \right)$$

$$= \left[\frac{1}{4} \left(1 - \cos(u) \right) \right]$$

(b) [4 points] $\iint_R x \, dA$ where R is the region between the circles $x^2 + y^2 = 1$ and $x^2 + y^2 = 4$ above the line y = 0 and below the line y = x.



$$\iint_{R} x \, dA = \int_{0}^{\pi/4} \int_{1}^{2} (r \cos 6) \, r \, dr \, d\theta = \int_{0}^{\pi/4} \cos 6 \cdot \int_{1}^{2} r^{2} \, dr \, d\theta$$

$$= \int_{0}^{\pi/4} \cos 6 \, \left(\frac{1}{3}r^{3}\right)_{r=1}^{r=2} \, d\theta = \int_{0}^{\pi/4} \cos 6 \, \left(\frac{1}{3}\cdot 8 - \frac{1}{3}\cdot 4\right) \, d\theta = \frac{7}{3} \int_{0}^{\pi/4} \cos 6 \, d\theta$$

$$= \frac{7}{3} \left(\sin 6\right)_{0}^{\pi/4} = \frac{7}{3} \left(\sin (\pi/4) - \sin (6)\right) = \frac{7}{3} \cdot \frac{\pi}{2} = \frac{7\sqrt{2}}{6}$$

(c) [2 points]
$$\int_{-\infty}^{\infty} e^{-2x^2} dx$$

$$I = \int_{-\infty}^{\infty} e^{-2x^2} dx$$

$$I^2 = \left(\int_{-\infty}^{\infty} e^{-2x^2} dx \right) \left(\int_{-\infty}^{\infty} e^{-2y^2} dy \right)$$

$$= \iint_{\mathbb{R}^2} e^{-2(x^2+y^2)} dA$$

$$= \int_{0}^{2\pi} \int_{0}^{\infty} e^{-2r^{2}} r dr d\theta \qquad u = 2r^{2} du = 4r dr$$

$$= \frac{1}{4} \int_{0}^{2\pi} \int_{0}^{\infty} e^{-2r^{2}} -4r dr d\theta$$

$$= \frac{1}{4} \int_{0}^{2\pi} \int_{0}^{\infty} e^{-u} du d\theta = \frac{1}{4} \int_{0}^{2\pi} (e^{-u})_{0}^{\infty} d\theta$$

$$= \frac{1}{4} \int_{0}^{2\pi} (-\frac{1}{2})_{0}^{\infty} - (-e^{0}) d\theta = \frac{1}{4} \int_{0}^{2\pi} d\theta = \frac{2\pi}{4}$$

$$= \frac{1}{4} \int_{0}^{2\pi} (-\frac{1}{4})_{0}^{\infty} - (-e^{0}) d\theta = \frac{1}{4} \int_{0}^{2\pi} d\theta = \frac{2\pi}{4}$$

$$= \frac{1}{4} \int_{0}^{2\pi} (-\frac{1}{4})_{0}^{\infty} - (-e^{0}) d\theta = \frac{1}{4} \int_{0}^{2\pi} d\theta = \frac{2\pi}{4}$$